

THEORETICAL PROBLEM No. 1
EVOLUTION OF THE EARTH-MOON SYSTEM
SOLUTIONS

1. Conservation of Angular Momentum

1a	$L_1 = I_E \omega_{E1} + I_{M1} \omega_{M1}$	0.2
1b	$L_2 = I_E \omega_2 + I_{M2} \omega_2$	0.2
1c	$I_E \omega_{E1} + I_{M1} \omega_{M1} = I_{M2} \omega_2 = L_1$	0.3

2. Final Separation and Angular Frequency of the Earth-Moon System.

2a	$\omega_2^2 D_2^3 = GM_E$	0.2
2b	$D_2 = \frac{L_1^2}{GM_E M_M^2}$	0.5
2c	$\omega_2 = \frac{G^2 M_E^2 M_M^3}{L_1^3}$	0.5
2d	The moment of inertia of the Earth will be the addition of the moment of inertia of a sphere with radius r_o and density ρ_o and of a sphere with radius r_i and density $\rho_i - \rho_o$: $I_E = \frac{2}{5} \frac{4\pi}{3} [r_o^5 \rho_o + r_i^5 (\rho_i - \rho_o)] .$	0.5
2e	$I_E = \frac{2}{5} \frac{4\pi}{3} [r_o^5 \rho_o + r_i^5 (\rho_i - \rho_o)] = 8.0 \times 10^{37} \text{ kg m}^2$	0.2
2f	$L_1 = I_E \omega_{E1} + I_{M1} \omega_{M1} = 3.4 \times 10^{34} \text{ kg m}^2 \text{ s}^{-1}$	0.2
2g	$D_2 = 5.4 \times 10^8 \text{ m}$, that is $D_2 = 1.4 D_1$	0.3
2h	$\omega_2 = 1.6 \times 10^{-6} \text{ s}^{-1}$, that is, a period of 46 days.	0.3
2i	Since $I_E \omega_2 = 1.3 \times 10^{32} \text{ kg m}^2 \text{ s}^{-1}$ and $I_{M2} \omega_2 = 3.4 \times 10^{34} \text{ kg m}^2 \text{ s}^{-1}$, the approximation is justified since the final angular momentum of the Earth is 1/260 of that of the Moon.	0.2

3. How much is the Moon receding per year?

3a	Using the law of cosines, the magnitude of the force produced by the mass m closest to the Moon will be: $F_c = \frac{G m M_M}{D_1^2 + r_o^2 - 2 D_1 r_o \cos(\theta)}$	0.4
3b	Using the law of cosines, the magnitude of the force produced by the mass m farthest to the Moon will be: $F_f = \frac{G m M_M}{D_1^2 + r_o^2 + 2 D_1 r_o \cos(\theta)}$	0.4
3c	Using the law of sines, the torque will be $\tau_c = F_c \frac{\sin(\theta) r_o D_1}{[D_1^2 + r_o^2 - 2 D_1 r_o \cos(\theta)]^{1/2}} = \frac{G m M_M \sin(\theta) r_o D_1}{[D_1^2 + r_o^2 - 2 D_1 r_o \cos(\theta)]^{3/2}}$	0.4
3d	Using the law of sines, the torque will be $\tau_f = F_f \frac{\sin(\theta) r_o D_1}{[D_1^2 + r_o^2 + 2 D_1 r_o \cos(\theta)]^{1/2}} = \frac{G m M_M \sin(\theta) r_o D_1}{[D_1^2 + r_o^2 + 2 D_1 r_o \cos(\theta)]^{3/2}}$	0.4
3e	$\tau_c - \tau_f = G m M_M \sin(\theta) r_o D_1^{-2} \left(1 - \frac{3r_o^2}{2D_1^2} + \frac{3r_o \cos(\theta)}{D_1} - 1 + \frac{3r_o^2}{2D_1^2} + \frac{3r_o \cos(\theta)}{D_1}\right)$ $= \frac{6 G m M_M r_o^2 \sin(\theta) \cos(\theta)}{D_1^3}$	1.0
3f	$\tau = \frac{6 G m M_M r_o^2 \sin(\theta) \cos(\theta)}{D_1^3} = 4.1 \times 10^{16} \text{ N m}$	0.5
3g	Since $\omega_{M1}^2 D_1^3 = G M_E$, we have that the angular momentum of the Moon is $I_{M1} \omega_{M1} = M_M D_1^2 \left[\frac{G M_E}{D_1^3} \right]^{1/2} = M_M [D_1 G M_E]^{1/2}$ The torque will be: $\tau = \frac{M_M [G M_E]^{1/2} \Delta(D_1^{1/2})}{\Delta t} = \frac{M_M [G M_E]^{1/2} \Delta D_1}{2 [D_1]^{1/2} \Delta t}$ So, we have that $\Delta D_1 = \frac{2 \tau \Delta t}{M_M} \left[\frac{D_1}{G M_E} \right]^{1/2}$ That for $\Delta t = 3.1 \times 10^7 \text{ s} = 1 \text{ year}$, gives $\Delta D_1 = 0.034 \text{ m}$. This is the yearly increase in the Earth-Moon distance.	1.0

3h	<p>We now use that</p> $\tau = -\frac{I_E \Delta \omega_{E1}}{\Delta t}$ <p>from where we get</p> $\Delta \omega_{E1} = -\frac{\tau \Delta t}{I_E}$ <p>that for $\Delta t = 3.1 \times 10^7$ s = 1 year gives</p> $\Delta \omega_{E1} = -1.6 \times 10^{-14} \text{ s}^{-1}.$ <p>If P_E is the period of time considered, we have that:</p> $\frac{\Delta P_E}{P_E} = -\frac{\Delta \omega_{E1}}{\omega_E}$ <p>since $P_E = 1 \text{ day} = 8.64 \times 10^4$ s, we get</p> $\Delta P_E = 1.9 \times 10^{-5} \text{ s}.$ <p>This is the amount of time that the day lengthens in a year.</p>	1.0
----	---	-----

4. Where is the energy going?

4a	<p>The present total (rotational plus gravitational) energy of the system is:</p> $E = \frac{1}{2} I_E \omega_{E1}^2 + \frac{1}{2} I_M \omega_{M1}^2 - \frac{G M_E M_M}{D_1}.$ <p>Using that</p> $\omega_{M1}^2 D_1^3 = G M_E, \text{ we get}$ $E = \frac{1}{2} I_E \omega_{E1}^2 - \frac{1}{2} \frac{G M_E M_M}{D_1}$	0.4
4b	$\Delta E = I_E \omega_{E1} \Delta \omega_{E1} + \frac{1}{2} \frac{G M_E M_M}{D_1^2} \Delta D_1, \text{ that gives}$ $\Delta E = -9.0 \times 10^{19} \text{ J}$	0.4
4c	$M_{\text{water}} = 4\pi r_o^2 \times h \times \rho_{\text{water}} \text{ kg} = 2.6 \times 10^{17} \text{ kg}.$	0.2
4d	$\Delta E_{\text{water}} = -g M_{\text{water}} \times 0.5 \text{ m} \times 2 \text{ day}^{-1} \times 365 \text{ days} \times 0.1 = -9.3 \times 10^{19} \text{ J}.$ Then, the two energy estimates are comparable.	0.3